

hw6 , due: Tuesday, October 24

1. a) A curve $y = f(x)$, $a \leq x \leq b$, is rotated about the line $y = c$, where $c \geq f(x)$. Find a formula for the resulting surface area. b) Using part (a), find the area of the surface obtained by rotating the circle $x^2 + y^2 = r^2$ about the line $y = r$.
2. An observer is located at height h above the North pole of a sphere of radius r . Show that the observer can see a portion of the sphere with area $S = \frac{2\pi r^2 h}{r+h}$. (hint: draw a picture, use the formula for the surface area obtained by rotating a curve about an axis)
3. Sketch the mass distribution, find the center of mass, and indicate the CM on the sketch.
 - a) three point masses on the x -axis; $m_1 = 25, m_2 = 20, m_3 = 10, x_1 = -2, x_2 = 3, x_3 = 7$
 - b) four point masses in the xy -plane; $m_1 = 6, m_2 = 5, m_3 = 1, m_4 = 4, (x_1, y_1) = (1, -2), (x_2, y_2) = (3, 4), (x_3, y_3) = (-3, -7), (x_4, y_4) = (6, -1)$
4. A triangular plate with uniform density has vertices $(-1, 0), (1, 0), (0, 2)$ in the xy -plane. Find the center of mass.
5. Sketch the region bounded by the curves $y = 4 - x^2$, $y = 0$ and find the center of mass.
6. Let $f(t)$ be the pdf for the time it takes to drive to work, where t is measured in minutes. Express the following probabilities as integrals.
 - a) the probability that the drive lasts 15 minutes or less
 - b) the probability that the drive takes a half hour or more
7. Let X be a random variable with pdf $f(x) = \frac{3}{64}x\sqrt{16 - x^2}$ for $0 \leq x \leq 4$ and $f(x) = 0$ for all other x .
 - a) Sketch $f(x)$ for $-\infty < x < \infty$. b) Verify that $f(x)$ is a valid pdf. c) Find $\text{prob}(0 \leq X \leq 2)$.
8. a) Evaluate $(1 + \frac{1}{n})^n$ for $n = 1, 10, 10^2, 10^3, 10^4$ using a calculator or Maple.
 - b) Evaluate $L = \lim_{n \rightarrow \infty} (1 + \frac{1}{n})^n$ using mathematics. (hint: compute $\ln L$)
9. This exercise introduces the hyperbolic functions.
 - a) Define $\cosh x = \frac{e^x + e^{-x}}{2}$, $\sinh x = \frac{e^x - e^{-x}}{2}$. Show that $\cosh x$ is an even function (i.e. $f(-x) = f(x)$) and $\sinh x$ is an odd function (i.e. $f(-x) = -f(x)$).
 - b) Sketch $y = \cosh x$ and $y = \sinh x$ on the same plot for $-\infty < x < \infty$.
 - c) Show that $\cosh^2 x - \sinh^2 x = 1$.
 note: This result shows that the point $(X, Y) = (\cosh x, \sinh x)$ lies on the hyperbola $X^2 - Y^2 = 1$ in the XY -plane, and this is why $\cosh x, \sinh x$ are called hyperbolic trigonometric functions. The functions $\cos x, \sin x$ are sometimes called circular trigonometric functions because they satisfy the equation $\cos^2 x + \sin^2 x = 1$, which implies that the point $(X, Y) = (\cos x, \sin x)$ lies on the circle $X^2 + Y^2 = 1$. By analogy with $\cosh x = \frac{e^x + e^{-x}}{2}$, $\sinh x = \frac{e^x - e^{-x}}{2}$, it can be shown that $\cos x = \frac{e^{ix} + e^{-ix}}{2}$, $\sin x = \frac{e^{ix} - e^{-ix}}{2i}$, where $i = \sqrt{-1}$; we'll discuss this later in the course as a consequence of Euler's formula.
 - d) Find $\frac{d}{dx} \cosh x$, $\frac{d}{dx} \sinh x$.
 - e) Define $\tanh x = \frac{\sinh x}{\cosh x}$. Show that $\tanh x$ is an odd function. Find $\frac{d}{dx} \tanh x$.
 - f) Evaluate $\lim_{x \rightarrow \pm\infty} \tanh x$. Sketch the graph of $\tanh x$ for $-\infty < x < \infty$.